

INTRODUCTIONS!

• Introduction

- Books

- Schedule for today, what we hope to accomplish

- Selective Mutism Association — www.selectivemutism.org

- Dr.Aimee Kotrba's monthly newsletter — sign up at www.thrivingmindsbehavioralhealth.com

• Involvement of audience, how to ask questions, making this interactive

**DEFINITION OF SM (DSM-5)** 

- Specific anxiety disorder
- Consistent, ongoing failure to speak in specific social situations, especially school
- $\bullet\,$  Not due to a primary language disorder
- Other disorders (e.g., stuttering, autism) have been ruled out
- A relatively rare childhood disorder, affecting approximately 1% of children in elementary school settings
- Behavior is deliberate self-protection, not deliberate oppositionality

2



COMMONTRAITS

Mutism


Blank facial expression, freezing, poor eye contact

Difficulty responding and/or initiating nonverbally

Slow to respond

Bilingual

3



PHYSIOLOGICAL CHARACTERISTICS

- Why don't children with SM look anxious???
- More chronically over-aroused than children with social phobia alone
  - Higher levels of arousal at baseline in studies, not just when asked to engage with others

 Children with SM appear to modulate their anxiety <u>better</u> than children with social anxiety, thereby not <u>appearing</u> as anxious outwardly.

/

10

### SHYNESS VS. SELECTIVE MUTISM

#### **SHYNESS**

- Slow warm up period
- Can often respond with a nod or small
- Same demeanor everywhere – quiet and reserved



- Warm-up time
   MUCH longer than expected
- Cannot respond at all -may appear frozen
- Dual personality restrained at school and talkative at home

PREVALENCE STATISTICS

- Most recent stats show approximately 1%...and growing?
- 1.5-2.6 / I female / male Garcia et al (2004)
- 90% comorbid with Social Anxiety
  - But different higher rates of oppositionality, agoraphobia, and language problems

3

## WHERE DOES SM ORIGINATE?

- NO evidence of causal relationship to abuse, neglect, or trauma
- Genetic predisposition model (genetic loading)
- Biological indicators
  - Decreased threshold of excitability in amygdala
  - Amygdala reacts more and takes longer to return to normal




Mean anxiety ratings per group

7
6
5
4
3
2
1
0
SM \_\_speech demanding \_\_embarrassing \_\_neutral

Schwenck, C., Gensthaler, A., & Vogel, F. (2019)

**SM VS SOCIAL ANXIETY** 



Anxious

Anxious-Oppositional

Anxious-Communication Delayed

TYPES OF SELECTIVE MUTISM

16

13

COEXISTING PROBLEMS
CONT.

Language Based Learning Disorder or communication deficits

Including pragmatics, grammar, semantics, articulation, voice, and fluency produce shorter, linguistically simpler, and less detailed language than typically developing children (McInnes, Fung, Fliscenbaum, & Tannock, 2004)
possibly weaker auditory-verbal memory span (Kristenson & Oerbeck, 2006)
lower receptive language scores than age matched peers (Nowakowski et al., 2009)

May be:


Independent of SM
Precursor to SM
Precursor to SM
Arising from lack of experience communicating due to the social anxiety of SM

Preventive
Factors

- High expectations
- Lack of family accommodation
- Involved parent and school in intervention

- Social isolation
- Denial of an issue
- Family accommodation
- Accidental reinforcement
- Speaking for the child

14 17



15 18

# SAFETY BEHAVIORS

- Anxious people often engage in a range of behaviors to make themselves feel safer when they cannot avoid anxious situations
- These behaviors are attempts to neutralize feelings of anxiety
- · High rate of "coercive behaviors"
  - Child truly believes the only way to alleviate anxiety is parent rescuing



ACCIDENTAL REINFORCEMENT

WHEN SOMEONE SAYS,
"I TRIED USING
REINFORCEMENT AND IT DIDN'T
WORK"

• Behaviorism 101 = reinforce whatever you want to maintain or increase

• How do school personnel and therapists accidentally reinforce?

• How might this impact anxiety?

19

CONCEPTUALIZING
SELECTIVE MUTISM

Child is prompted to speak or engage

Increased likelihood of avoidance

Child avoidance

Child avoidance

Child avoidance

Child avoidance

Child avoidance

Child is rescued

I can't talk to talk but I just don't know how to say it.

If I talk I might say the wrong thing.

If I talk everyone will look at me!

23

20

AVOIDANCE — THE OXYGEN OF ANXIETY

KEETS KIDS FROM LEARNING THAT FEAR IS A WARNING, NOT TO THE WORLD

KEETS KIDS FROM LEARNING TO STEER THEMSELYES AWAY FROM UNFADOCTABLE OR UNCOMPORTABLE STUATIONS

SCHOOL IMPLICATIONS

- Academic implications
- Inability to assess skills (especially reading)
- Possibly limited peer relationships
- Behavioral implications (participation)
- Social Implications (how peers see child)

21 24

# Worsening anxiety and depression Social isolation, impaired peer relationships Self-esteem issues School refusal, poor academic performance Self-medication Problems adjusting to work situations

- Everyday that a child continues with impairing symptoms:
  - Is <u>not</u> neutral
  - Strengthens the habit of avoidance
  - Strengthens perceptions that they are the child that doesn't talk
  - Decreases self-confidence

Selective Mutism
 Questionnaire
 SCARED (Screen for
 Childhood Anxiety Related
 Disorders)
 And....
 Autism Diagnostic
 Observation Schedule
 (ADOS)
 Speech/language evaluation
 IQ (nonverbal)
 Etc...

25 28





I. Add totals in each 0+0+0+0+0+0=0 /6=School=0 (Prinfer Intent) of ordered.

3. When collect on by his or her issurber, my child sesson. Divide by number of items in section For total 3+2+2+3+2+2= |4/6= score, add up totals in each Family=2.33 section – DO NOT 0+1+2+0+0=3/ divide 5= Other=.6 Total 0+14+3=17

26 29

STIC INTER	VIEW
Family	Child
Genetic history     Home life     description     Recent     stressors	Behavioral characteristics     Medical history     Repetitive or restricted interests, obsessive thoughts, etc.
	Family  • Genetic history  • Home life description  • Recent





TYPES OF TREATMENT

34

Speech Frequency (How?)	Communication Partners (Who?)	Setting Variable (Where?)	Intervention Recommended
Limited	Most people	Most environments	Contingency management
Typical	One or limited people	Most environments	Stimulus fading of new people in specific environments
Typical	Most people	One environment	Stimulus fading of environments
Limited	One or limited people	One environment	Stimulus fading of new people into comfortable environment AND stimulus fading of environments
No speech	No people	No environments	Shaping AND Stimulus fading of new people and environments

**OVERVIEW OF BEHAVIORAL TREATMENT** • Emphasis placed on environmental determinants of  $\bullet\,$  Behavior was, to some degree, a result of learning, and can therefore be unlearned Treatment consists of modifying environment, systematically practicing new behaviors, and identifying factors that maintain avoidance behavior • Belief that kids do well if they can

32 35

# LITERATURE/RESEARCH ON SELECTIVE MUTISM

**RESEARCH ON BEHAVIORAL TREATMENT**  Most supported treatment per research Kratochwill, 1981; Krohn et al, 1992; Leonard & Topol, 1993; Tancer, 2002; Oerbeck et al, 2014; Bergman, 2014 Effective in increasing production of speech in social situations - Stone & Kratochwill, 2002 Even more effective when combined with behavioral school and home-based exposure program - Bergman, 2005 Exposure-based practice may be more effective than parent-focused contingency management - Vecchio & Kearney, 2008

33 36

# RESEARCH ON OTHER INTERVENTIONS

- Psychoanalysis, DIR/Floortime, play therapy
- No real experimental research published; some case studies.
- No evidence of effectiveness for treatment of Selective Mutism

## **MEDICATION**

#### UNLIKELY TO MEDICATE

- · Less severe impairment
- No CBT trial in the past
- Low comorbities

mutism

- · Family history not strong
- Meeting CBT benchmarks

### MORE LIKELY TO MEDICATE

- · More severe impairment
- Poor prior CBT response
- · High comorbidities
- · Strong family history
- Not meeting CBT benchmarks

37

40


# MEDICATION AS AN INTERVENTION?

- · Addresses biological determinants
- Most common medication utilized and researched for SM is Prozac (Fluoxetine), a Selective Serotonin Reuptake Inhibitor
- NOT effective alone
- · Goal is use as water wings
- Goal is usually to have the child take the medication for 9-12 months
- Concerns?



41

38



EVIDENCE-BASED INTERVENTION BEHAVIORAL TREATMENT

RESEARCH ON PSYCHOPHARMACOLOGICAL INTERVENTIONS

 Overall, research on Prozac suggests it is helpful in reducing anxiety and increasing social interactions. (Carlson, Mitchell, & Segool, 2008)


· Minimal support for Zoloft (Sertraline) as a beneficial treatment


· Currently, no medications have achieved FDA approval for the

treatment of childhood social phobia or SM.


 Two research studies support other SSRIs as beneficial medications. (Lehman, 2002;Thomsen, Rasmussen, & Anderson, 1999)

- Differences in parent report and teacher/clinician report of child's







**PSYCHO-EDUCATION FOR KIDS** • Explain in developmentally appropriate speech - Building brave muscles - Practicing to get better - Starts hard, gets easier - How our brains send the "emergency" signal even when it's not necessary – we can learn to ignore or overpower the signal. - Warheads example




44





45 48



## **GOAL OF TREATMENT**

- NOT making anxiety go away!
- Can't talk your amygdala out of a fear it's not listening! (signals between the cortex and amygdala run mostly one way)
- Our amygdala has to be activated to learn something new activate it, hang around and nothing terrible happens, amygdala learns.
- Learning to be "comfortable with discomfort!"
- To build distress tolerance (in <u>adults</u> AND the kids!) through facing fears.

49 52

# WHAT IS DESENSITIZATION?


- Increasing ability to communicate slowly through facing fears at a reasonable pace
- Stops pattern of reinforcement of avoidance
- Allows for slow decrease of anxiety
- Demonstrates successes, which increases motivation
- Practice, practice, practice!

**KEY PLAYERS -**

- · Desensitization is done by:
  - Psychologist/mental health professional outside school (helping with family training and public practice, consulting with school)
  - Parent
  - Keyworker school personnel who is primarily in charge of:
    - Desensitizing (exposing)
    - Generalizing to all school environments
    - Communicating with teacher, parent, psychologist



50 53





51 54



#### DOS

- · Labeled praise for talking
- Reflecting verbalizations
- Play-by-play announcer
- Forced choice/open ended questions
- Direct prompts to talk
- Wait 5 seconds


DON'TS

• Mind reading


• Yes/no questions

- Indirect commandsNegative talk
- Enabling/rescuing

55



58



## **STIMULUS FADING**

- Gradually increasing the number of different people the child speaks to and settings the child speaks in
  - Start where the child currently speaks
  - Gradually introducing new people into conversations
  - Speaking in new settings with the help of stimulus associated with speaking (e.g., communication ladders with speaking partner or keyworker)

**GENERALIZING SKILLS** 

- Stimulus fading video

Pathway HazardsPeople

-Environment

-Speech demands

\*\*Only change one at a time!

59

# FORCED-CHOICE QUESTIONS

- Forced choice questions increase the likelihood of a verbal response!
  - Practice developing forced-choice questions.
    - What are you going to be for Halloween?
    - Do you want a brownie?Do you have a sister?
    - What is your favorite school subject?
    - Do you remember where we are going on the field trip today?

#### **COMMUNICATION LADDER** (SHAPING)

- Shaping
  - $\boldsymbol{\mathsf{-}}\ \mathsf{Reinforcing}\ \mathsf{successive}\ \mathsf{approximations}\ \mathsf{of}$
  - Initially reinforcing more frequently occurring behaviors (nonverbals, simple sounds, etc.)
  - Gradually reinforcing behaviors that approximate full speech (words, sentences, etc.)

**EXAMPLES** 

#### SCAFFOLDING

- Answering to a parent 5 feet from counter at the ice cream store, then moving closer the next time to the parent
- Reading to small group with peer and teacher, then fading in more peers
- Systematic plan for show and tell- on video, then in front of small group, then class

#### ENABLING

- · Child orders to parent, 5 feet from counter, always
- Only reading in small group all year
- Use of iPad for Show and tell all year

61

64

### **COMMUNICATION LADDER** INDEPENDENT Spontaneous Initiating Multiple word, longer utterances Single words Sounds to indicate meanings Voiced sounds Voiceless sounds ) Blowing air Nonverbal communication HIGH SUPPORT Following directions

**TRACKING BRAVE PRACTICES** 

/	Date:	Location:	Communication Partner:	Content of Speech:	Notes:
$\langle$	10/22	SLP's Office	Speech Pathologist	Named colors in game with one-word responses to forced-choice questions	Quiet but audible speech. Responded well to rewards.
/	10/25	Hallway	Speech Pathologist	Did Favorite's Game in public location	Notable reduction in volume when others came by.
$\setminus$	10/31	SLP's Office	Speech Pathologist and Classroom Teacher	Played jenga game with questions while teacher faded in.	No reduction in volume; very brave!!

When to Increase to a harder demand

62

63

65

# **SCAFFOLDING VS. ENABLING/RESCUING**

#### SCAFFOLDING


- Understanding a child's current ability and asking them to push themselves consistently outside of their comfort zone
- Knowing what is TOO far and assisting them in bridging the gap to still be successful/participate

#### RESCUING

- Setting the bar at or below the child's current functioning
- Allowing avoidance to occur or continue
- Stepping in before the child has a chance to manage their own anxiety and make an effort


TIPS/FOR ENGOURAGING SPEECH

- REMAIN CALM!!!
- · Use specific praise
- Judicious use of direct prompts to speak
- Brave talking is target behavior, not correctness
- Always wait 5 seconds for reply child needs opportunity to respond
- Try to always ask forced-choice instead of yes/no questions (to avoid headshaking)
- · Use situations that are motivationally driven to encourage more speech
- Don't mind read





67 70



DOS AND DON'TS OF **INTERVENTION** DON'T.... Allow avoidance to continue unchecked Work with school to create intervention plan • Rescue · Advocate for child Plateau · Be creative with rewards. Forget about both responding AND initiating practices, goals, etc. Use stimulus associated with Say "don't worry – no one will hear you!" speech (you!) to elicit speech in school • Quit! © Identify avoidance and work toward approach or scaffolding

68 71

Perhaps intensive doses are better
 Pros: Less warm-up time, likely quicker outcomes, avoiding negative outcomes of continuing behavior, return to appropriate developmental tasks, greater access to treatment, cost effectiveness
 Cons: cost up front, not covered by insurance, exhausting, interrupts daily activities

COMMUNITY

69 72

# **CREATING A**

- Determine appropriate community practices: Child's anxiety level should always be the guide Fit practices into ordinary outings

- Preparing your child

  - Discuss why you're practicing
    Quantifying anxiety (e.g., I-5 scale, red/yellow/green light)
  - Consider environmental variables (e.g., time of day)
  - Discuss motivational factors
  - Rehearse several times
- Preparing the community person

   Simple explanation of the purpose
- Prompt the person with what you'd like to say (script it exactly!)
- Don't be afraid to correct!





# **DEALING WITH WELL-**

- · Community persons often make well-meaning mistakes:
  - $\,-\,$  Looking to the adult to speak for the child
  - Allow nonverbals
  - Encouraging nonverbals ("You can point to your choice")
  - Asking you what the child said



- Prepare the person in advance (if possible)
- Look at the child
- Politely give them a script of what to say
- Remember, you won't likely see this person again so if its uncomfortable, its

73



# TRACKING COMMUNITY **BRAVE PRACTICES**

Date:	Location:	Communication Partner:	Content of Speech:	Notes:
10/22	Red Robin	Server	Responded drink and food choices in one word answers ("Cheeseburger")	Loud environment; speech was audible
10/25	Target	3 Employees	Mom said "Where can we find" and Lauren gave name of items	Speech became louder with each practice
10/31	Local	Neighbors	Said "Trick or Treat" with siblings at several homes	Visibly less anxious after each practice

When to Increase to a harder demand

## **FAMILY MEMBERS**

- Education
- Practice Partners
- · Additional advocates in community settings



74

77

## **EXAMPLES OF SCAFFOLDING V. RESCUING IN COMMUNITY**

#### SCAFFOLDING

- Community person can't hear child, parent/employee asks child to speak up, the child speaks louder and parent amplifies speech
- Child says he isn't able to do the practice, parents help child find a Plan B that is somewhat easier
- Child nonverbally indicates ice cream choice, parents script a question for the employee to ask


#### RESCUING


- Community person can't hear child so the parent takes over and answers all additional questions
- Child tells parents he isn't able to practice, parents allow child to avoid that practice
- · Child nonverbally indicates ice cream choice and gets ice cream

#### **EXTRACURRICULAR ACTIVITIES**

- Consider sharing some information
  - Quick facts regarding SM and ways to engage a child with SM
- · Arrive early
- · Playdates with teammates
- · Visit the location periodically







79 82





80

# STIMULUS FADING ONLINE

- Parent and child play a game with computer/phone/ipad nearby
- New person signs on and busies themselves in background
- New person moves closer to screen but ignores
- $\bullet\,$  New person begins to comment on the activity, then reflects what child is saying
- $\bullet\,$  New person relays questions through parent and then reflects
- $\bullet\,$  New person asks FC questions directly of the child
- Parent begins to disengage
- Parent leaves the room or goes as far away as possible



81 84



SPECIFIC SCHOOL INTERVENTIONS AND ACCOMMODATIONS

- Desensitization in school with keyworker
  - 5-10 minutes of practice daily
  - As structured as possible
  - Consider fading with teacher and PW
  - · Team meetings and communication
- Participating (at least) nonverbally
- · Chores and responsibilities
- Extracurriculars
- Seating arrangement/small group activities
- Conversational partners/conversational visits
- Allow for early arrival (warm-up time)

88

IEP or Section 504? Which is better?
How do I request a special education plan?
If IEP, what identification?
Other Health Impaired - Limited strength, vitality or alertness due to chronic or acute health problems which adversely affects educational performance - 34 C.F.R. § 300.8(a)(9)

Speech/Language Impaired - Communication disorder, such as

 Speech/Language Impaired - Communication disorder, such as suttering, impaired articulation, a language impairment, or a voice impairment that adversely affects educational performance - 34 C.F.R. § 300.8(a)(11)

 Emotional Disturbance - Condition exhibiting one or more of the following characteristics over a long period of time and to a marked degree that adversely affects educational performance -34 C.F.R. § 300.8(a)(4) SPECIFIC SCHOOL INTERVENTIONS,

- Nonverbal evaluation methods
  - Audio or videotaping
  - Writton tosting
  - Parent report
- Provide practice experience in advance
- Transition plan for the next year begins in April
- Teacher and peer selection for class
- Do school visits/teacher visit during summer

Scheduled times/bathroom buddy for bathroom breaks

86 89

# WHICH IS BETTER - IEP OR SECTION 504?

IEP

- Disability that adversely affects educational performance
- Child requires special education and direct instruction
- Child may require related services

#### SECTION 504

- Currently have a disability which affects a major life activity
- Child requires accommodations to access major life activities
- Child may require related services

#### FACTORS TO CONSIDER IN CREATING THE RIGHT ENVIRONMENT FOR SM KIDS

- Familiarity of audience (but beware "contamination"!)
- Gender
- Size of audience
- Location (private vs. public)
- Expectation of response (open ended vs. yes/no vs. forced choice)
- Eye-contact
- Language purpose tested or graded?

87 90



- By a few sessions, child should not look angry or frightened to start sessions
- By a few sessions, progress should be obvious - even if slow but steady
- By 2-3 sessions, children are usually talking to me in the room alone
- · By 4-6 sessions, children are usually talking to me with parents/siblings in



Minimizes negative impact on the Prevents situation from becoming **IMPORTANCE** Prevents mutism from becoming OF EARLY engrained INTERVENTION Prevents repeated ineffective attempts to elicit speech Minimizes emotional and physical strain caused to parents and teachers

91 94


By 6-8 in-school weeks, most are talking to an adult without parent present in room

By 8-12 in-school weeks, most are talking to multiples of teachers or peers, even if in contrived situations

By 12 weeks the child should seen confident that gains are being made

GENERAL TREATMENT **BENCHMARKS** 






95

92

**TREATMENT** REVIEW

**OUESTIONS?** 

Thriving Minds - $\underline{www.thriving minds behavioral health.com}$ (734) 433-5100 akotrba@thrivingminds.info



93 96